

CLAIMS

1. A diagnostic drug for autoimmune diseases, comprising at least one of a polypeptide selected from an HMG-1 family, a polypeptide selected from an HMG-2 family, a fragment thereof which is reactable with an antibody of an autoimmune disease patient,

wherein when the polypeptide is a neutrophil 28kDA antigen, the autoimmune disease is not ulcerative colitis.

2. A diagnostic drug for autoimmune diseases according to claim 1, wherein the autoimmune disease is rheumatoid arthritis, systemic lupus erythematosus, Sjögren's syndrome, Behçet's disease, scleroderma, primary biliary cirrhosis, microscopic polyangiitis/polyarteritis nodosa, ulcerative colitis, Crohn's disease, or autoimmune hepatitis.

3. A diagnostic drug for autoimmune diseases according to claim 1, wherein the polypeptide is selected from human, bovine, porcine, chicken, mouse or rat HMG-1 or HMG-2.

4. A kit for diagnosing autoimmune diseases, comprising at least one of a polypeptide selected from an HMG-1 family, a polypeptide selected from an HMG-2 family, a fragment thereof which is reactable with an antibody of an autoimmune disease patient,

wherein when the polypeptide is neutrophil 28kDA, the autoimmune disease is not ulcerative colitis.

5. A kit for diagnosing for autoimmune diseases according to claim 4, wherein the autoimmune disease is rheumatoid arthritis, systemic lupus erythematosus, Sjögren's

09214881.050799

Sub  
a'

syndrome, Behçet's disease, scleroderma, primary biliary cirrhosis, microscopic polyangiitis/polyarteritis nodosa, ulcerative colitis, Crohn's disease, or autoimmune hepatitis.

6. A kit for diagnosing for autoimmune diseases according to claim 4, wherein the polypeptide is selected from human, bovine, porcine, chicken, mouse or rat HMG-1 or HMG-2.

7. A method for detecting an antibody of autoimmune disease patient, comprising the step of reacting a reagent including at least one of a polypeptide selected from an HMG-1 family, a polypeptide selected from an HMG-2 family, a fragment thereof which is reactable with an antibody of an autoimmune disease patient,

wherein when the polypeptide is a neutrophil 28kDA antigen, the autoimmune disease is not ulcerative colitis.

8. A method according to claim 7, wherein the autoimmune disease is rheumatoid arthritis, systemic lupus erythematosus, Sjögren's syndrome, Behçet's disease, scleroderma, primary biliary cirrhosis, microscopic polyangiitis/polyarteritis nodosa, ulcerative colitis, Crohn's disease, or autoimmune hepatitis.

9. A method according to claim 7, wherein the polypeptide is selected from human, bovine, porcine, chicken, mouse or rat HMG-1 or HMG-2.

Sub A2

Sub A3

Add A4

Add H2

06214831-060799